

Intracystic Injection of Large Surface Area Microparticle Paclitaxel for Chemoablation of Intraductal Papillary Mucinous Neoplasms: Insights from an Expanded Access Protocol

Devarshi R Ardeshta, Zarine K Shah, Phil A Hart, Georgios Papachristou, Samuel Han, Peter Lee, Mary Dillhoff, Andrei Manilchuk, Jordan Cloyd, Aslam Ejaz, Anne Mary Noonan, Timothy M Pawlik, Megan E Fry, Somashekar G Krishna

Division of Gastroenterology, Hepatology, and Nutrition, The Ohio State University Wexner Medical Center, Columbus, Ohio, 43210 USA



THE OHIO STATE UNIVERSITY
WEXNER MEDICAL CENTER

BACKGROUND and AIMS

Branch duct (BD) – Intraductal Papillary Mucinous Neoplasms (IPMNs) represent 80% of incidentally identified pancreatic cystic lesions (PCLs). Guideline-directed risk assessment is suboptimal and about 60% patients undergo unnecessary surgical resection. Thus, minimally invasive treatments are increasingly being utilized.

A phase 1 clinical trial evaluated the safety, tolerability, and preliminary efficacy of EUS-guided fine needle injection (EUS-FNI) with novel particle-engineered form of paclitaxel with large surface area microparticles (LSAM-PTX, NanoPac®, CritiTech, Lawrence, KS) in mucinous PCLs.

We prospectively evaluated the safety and response of EUS-FNI of LSAM-PTX to chemoablate BD-IPMNs in an expanded access protocol (EAP).

METHODS

Subjects diagnosed with BD-IPMNs and deemed non-surgical received up to two doses of LSAM-PTX (15mg/mL; 12 weeks apart) by EUS-FNI as part of a multicenter clinical trial (NCT03188991). Subjects at our center were subsequently enrolled into an EAP where additional doses of LSAM-PTX were administered by EUS-FNI on similar schedules (Figure 1).

RESULTS

Six BD-IPMNs measuring (mean ± standard deviation (SD)) 3.18 ± 0.76 cm in diameter among 5 subjects (mean age: 66 years) were treated by EUS-FNI of LSAM-PTX. Two to six doses of LSAM-PTX (mean dose/cyst: 73 ± 31 mg) were administered, and subjects were followed for up to 32 months.




Figure 1. Study flow chart

The mean volume reduction of the treated cysts was 42-89% (absolute volume change from 9.58 ± 5.1ml to 2.2 ± 1.1ml (p=0.016)), Figure 2. The mean surface area was reduced from 31-83% (absolute reduction from 21.9 ± 8.7cm² to 5.7 ± 2.5cm² (p=0.009)), Figure 2.

Higher dosing-frequency of EUS-FNI of LSAM-PTX significantly correlated with a reduction in cyst volume (R²=0.87, p=0.03) and surface area (R²=0.83, p=0.04), Figure 3.

Comparing pre-and-post ablation, molecular analysis of the cyst fluid demonstrated loss of the cyst-associated mutations in 5 (83.3%) with reemergence in 1 and persistence in 1, and intracystic changes of fibrosis or calcification were observed in 83.3% (n=5) BD-IPMNs (Figure 4).

Immediate and Sustained Response




Figure 2. Changes in BD-IPMN volume and surface comparing pre-ablation, immediate and sustained response; box-plot with 95% CI.




Figure 3. Reduction in BD-IPMN volume and surface and correlation with frequency and dosing of LSAM-PTX

ADVERSE EVENTS

One subject developed mild acute pancreatitis following EUS-FNI (5th dose) of LSAM-PTX in a fibrotic BD-IPMN.

	Phase	CT and/or MRI changes	EUS morphology	Cyst fluid molecular analysis
BD-IPMN #1	Pre-ablation	Septations, septal and wall enhancement, focal dilation of PD	Multilobulated, thick wall and septations	KRAS (p.G12V) GNAS (p.R201H)
	Post-ablation	Small to characterize, fibrosis, decrease in focal dilation of PD	Increase in size, inflammatory debris no septations, thick wall	No mutations
	Sustained			No mutations
BD-IPMN #2	Pre-ablation	Septations, septal enhancement	Multilobulated, thin septations, thin wall	KRAS (p.G12V) GNAS (p.R201C)
	Post-ablation	New calcifications, decrease in size	Reduced size, thick septations, thick wall, fibrosis	KRAS (p.G12V) GNAS (p.R201C)
	Sustained			
BD-IPMN #3	Pre-ablation	Uniform thin wall, no septations	Single compartment, thin wall	GNAS (p.R202C) BRAF (p.V600_S605delinsD)
	Post-ablation	Decrease in size, new septations	Decrease in size, subtle septations vs. fibrosis	No mutations
	Sustained			
BD-IPMN #4	Pre-ablation	Dense septations, thin wall	Multilobulated, thin septations, thin wall	KRAS (p.Q61H) GNAS (p.R202C)
	Post-ablation	Prominent septations, fibrotic changes, decrease in size	Reduced size, multilobulated, thick septations, thick wall	KRAS (p.Q61H) GNAS (p.R202C)
	Sustained			
BD-IPMN #5	Pre-ablation	Lobulated, thin septations, thin wall	Multilobulated, thin septations, thin wall	KRAS (p.G12V) GNAS (p.R202H)
	Post-ablation	Decrease in size, thick septations, fibrotic changes	Reduced size, multilobulated, thin septations, thin wall	No mutations
	Sustained			
BD-IPMN #6	Pre-ablation	Enhancement of smooth cyst wall	Single compartment, thick wall	KRAS, p.G12R
	Post-ablation	Thick wall, fibrosis, calcification	Inflammatory debris, thickening of wall	No mutations
	Sustained			

Figure 4. Morphological changes observed in cross-sectional imaging and EUS after EUS-guided chemoablation with LSAM-PTX

CONCLUSION

In this extended access protocol, EUS-FNI of LSAM-PTX into BD-IPMNs was safe and resulted in volume and surface area reduction, morphological changes, and loss of pathogenic mutations.